

Livestock

Corn feed meal PL

Corn feed meal PL

is released during the processing of maize into maize semolina meal and maize flour for the food industry. Corn feed meal PL consist of bran, germs and parts of the mealy core. Corn feed meal PL can be used perfectly as a replacement for CCM.

Quality

The product complies with laws and regulations. Mycotoxins, insecticides, pesticides and herbicides are tested according to the GMP+ FSA. The product is non-GMO (according to Regulation (EC) 1829/2003 and 1830/2003).



Valid from 08 November 2022

Downloaded on 15 December 2025

		AID	SID
Dry matter %	86.4	2.2	2.6
pH	6.4	1.9	2.0
Crude protein	105	3.6	3.9
Crude fat	102	2.4	3.0
Crude fibre	45	0.4	0.5
Crude ash	28	2.4	2.8
Starch total	539	8.4	8.9
Sugar	33	3.6	4.1
NSP	226		
Calcium (Ca)	0.7		1.47
Sodium (Na)	0.0		3101
Chloride (Cl)	0.6		16.2
Potassium (K)	7.2		
Magnesium (Mg)	2.1		
Phosphorus total (P)	6.0		
Dig. Phosphorus	1.5		
Dig. Lysine			
Dig. Methionine			
Dig. Methionine + Cystine			
Dig. Threonine			
Dig. Tryptophan			
Dig. Isoleucine			
Dig. Leucine			
Dig. Valine			
Energie value			
NE pigs, kcal			
ME Schw., MJ			
Cattle values			
DVE			111
OEB			-62
nXP			175.85
Starch resistant			118
FOS			586
VEM			1261
VEVI			1400
NEL			8.98
ME Rind, MJ			14.20

Levels are based on averages and stated in grams per kg of dry matter, unless stated otherwise. Changes in product composition are reserved.

Feeding advice based on DS of total ration

Weaned piglets (up to 25 kg)	max 20%
Fattening pigs (25 - 50 kg)	max 25%
Fattening pigs (from 50 kg)	max 30%
Sows	max 30%
Dairy cattle	max 4 kg product in ration
Beef cattle	max 1 kg product / 100 kg body weight

Logistics & Storage

Transport	Bulk with tipper, walking floor or blown in silo
Storage	Store in a dry, dark and cool place or in a closed dry feed silo. Clean storage areas regularly.
Shelf life	3 months
Nipple worthy	No