

Livestock

Corn feed meal PL

Corn feed meal PL

is released during the processing of maize into maize semolina meal and maize flour for the food industry. Corn feed meal PL consist of bran, germs and parts of the mealy core. Corn feed meal PL can be used perfectly as a replacement for CCM.

Quality

The product complies with laws and regulations. Mycotoxins, insecticides, pesticides and herbicides are tested according to the GMP+ FSA. The product is non-GMO (according to Regulation (EC) 1829/2003 and 1830/2003).



Valid from 08 November 2022

Downloaded on 02 February 2026

Dry matter %	86.4
pH	6.4
Crude protein	105
Crude fat	102
Crude fibre	45
Crude ash	28
Starch total	539
Sugar	33
NSP	226
Calcium (Ca)	0.7
Sodium (Na)	0.0
Chloride (Cl)	0.6
Potassium (K)	7.2
Magnesium (Mg)	2.1
Phosphorus total (P)	6.0
Dig. Phosphorus	1.5

	AID	SID
Dig. Lysine	2.2	2.6
Dig. Methionine	1.9	2.0
Dig. Methionine + Cystine	3.6	3.9
Dig. Threonine	2.4	3.0
Dig. Tryptophan	0.4	0.5
Dig. Isoleucine	2.4	2.8
Dig. Leucine	8.4	8.9
Dig. Valine	3.6	4.1
 Energie value		
1.47		
NE pigs, kcal		
3101		
ME Schw., MJ		
16.2		

Cattle values

DVE	111
OEB	-62
nXP	175.85
Starch resistant	118
FOS	586
 VEM	
1261	
VEVI	1400
NEL	8.98
ME Rind, MJ	14.20

Levels are based on averages and stated in grams per kg of dry matter, unless stated otherwise. Changes in product composition are reserved.

Feeding advice based on DS of total ration

Weaned piglets (up to 25 kg)	max 20%
Fattening pigs (25 - 50 kg)	max 25%
Fattening pigs (from 50 kg)	max 30%
Sows	max 30%
Dairy cattle	max 4 kg product in ration
Beef cattle	max 1 kg product / 100 kg body weight

Logistics & Storage

Transport	Bulk with tipper, walking floor or blown in silo
Storage	Store in a dry, dark and cool place or in a closed dry feed silo. Clean storage areas regularly.
Shelf life	3 months
Nipple worthy	No