

Livestock

Wheat starch Tarpro

Wheat Starch Tarpro is released during the extraction of starch from wheat, resulting in several valuable liquid products for livestock farming. Tarpro, liquid wheat starch, is one of these products. The whole wheat grain is cleaned, conditioned, ground and sieved, after which wheat flour, wheat germ and wheat bran are released. The wheat flour is separated into wheat starch and wheat protein (gluten) using water and physical processes. Co-products are released from the various processes, including a starch-rich and a protein-rich flow that form the product Tarpro. Tarpro is thickened and acidified.

Quality

The product complies with laws and regulations. Mycotoxins, insecticides, pesticides and herbicides are tested according to the GMP+ FSA. The product is non-GMO (according to Regulation (EC) 1829/2003 and 1830/2003).



Valid from 22 March 2023

Downloaded on 02 February 2026

Dry matter %	24.0
pH	3.8
Crude protein	143
Crude fat	33
Crude fibre	3
Crude ash	49
Starch total	306
Sugar	427
Calcium (Ca)	1.2
Sodium (Na)	5.7
Chloride (Cl)	4.0
Potassium (K)	10.9
Phosphorus total (P)	4.4
Dig. Phosphorus	2.7
Sulfate	17.9
Lactic acid	13
Acetic acid	1
Alcohol	4

	AID	SID
Dig. Lysine	5.7	6.1
Dig. Methionine	2.1	2.2
Dig. Methionine + Cystine	4.9	5.3
Dig. Threonine	4.0	4.5
Dig. Tryptophan	1.3	1.4
Dig. Isoleucine	3.8	4.2
Dig. Leucine	7.5	8.0
Dig. Valine	6.4	6.9
Energie value		
1.46		
NE pigs, kcal		
3062		
ME Schw., MJ		
16.2		

Levels are based on averages and stated in grams per kg of dry matter, unless stated otherwise. Changes in product composition are reserved.

Feeding advice based on DS of total ration

Weaned piglets (up to 25 kg)	max 10%
Fattening pigs (25 - 50 kg)	max 15%
Fattening pigs (from 50 kg)	max 20%
Sows	max 20%

Logistics & Storage

Transport	Liquid with tank truck
Storage	In acid-resistant silo or bunker with agitator, regular stirring is desirable to prevent settling.
Shelf life	3 months, provided the silo or bunker is cleaned regularly
Nipple worthy	Yes