

Livestock

Brewer's yeast

Brewer's yeast is produced when brewing beer. Much brewer's yeast is processed into yeast products for human use. In addition, brewer's yeast in dried and liquid form is sold as animal feed. Brewer's yeast contains a lot of protein with an excellent amino acid pattern. Animal's health can be positively influenced by the presence of B vitamins. Brewer's yeast contains between 5 - 7% of alcohol.

Quality

The product complies with laws and regulations. Mycotoxins, insecticides, pesticides and herbicides are tested according to the GMP+ FSA. The product is non-GMO (according to Regulation (EC) 1829/2003 and 1830/2003).



Valid from 21 December 2022

Downloaded on 02 February 2026

| | |
|----------------------|------|
| Dry matter % | 10.5 |
| pH | 4.8 |
| Crude protein | 443 |
| Crude fat | 25 |
| Crude fibre | 10 |
| Crude ash | 77 |
| Starch total | 182 |
| Sugar | 26 |
| NSP | 283 |
| Calcium (Ca) | 1.3 |
| Sodium (Na) | 0.9 |
| Chloride (Cl) | 1.6 |
| Potassium (K) | 17.7 |
| Magnesium (Mg) | 1.4 |
| Phosphorus total (P) | 11.6 |
| Dig. Phosphorus | 5.8 |
| Lactic acid | 20 |
| Acetic acid | 6 |
| Alcohol | 452 |



Loop Co-Products BV

Parlevinklerweg 8
5928 NV Venlo
The Netherlands

www.loop.company
support@loop.company
+31(0)88 56 66 721

IBAN:

BIC:

VAT. NL

VAT. BE

UST.ID:

C.C. Venlo:

NL55 RABO 0321 8217 85

RABO NL2U

NL0095.32.488.13.01

BE0547739105

DE2891.81.759

1203 5225



GMP+:

QS ID:

GMP018172

4031735219906

| | AID | SID |
|---------------------------|-----|------|
| Dig. Lysine | 0.0 | 26.1 |
| Dig. Methionine | 0.0 | 5.7 |
| Dig. Methionine + Cystine | 0.0 | 9.1 |
| Dig. Threonine | 0.0 | 17.6 |
| Dig. Tryptophan | 0.0 | 4.5 |
| Dig. Isoleucine | 0.0 | 17.1 |
| Dig. Leucine | 0.0 | 25.6 |
| Dig. Valine | 0.0 | 19.7 |
| Energie value | | |
| 2.14 | | |
| NE pigs, kcal | | |
| 4502 | | |
| ME Schw., MJ | | |
| 22.9 | | |

Levels are based on averages and stated in grams per kg of dry matter, unless stated otherwise. Changes in product composition are reserved.

Feeding advice based on DS of total ration

| | |
|------------------------------|---------|
| Weaned piglets (up to 25 kg) | max 5% |
| Fattening pigs (25 - 50 kg) | max 8% |
| Fattening pigs (from 50 kg) | max 12% |
| Sows | max 10% |

Logistics & Storage

| | |
|---------------|---|
| Transport | Liquid with tank truck |
| Storage | In acid-resistant silo or bunker with agitator, regular (quiet) stirring is desirable |
| Shelf life | 2 weeks, provided the silo or bunker is cleaned regularly. Acidification can extend the shelf life up to 6 weeks. |
| Nipple worthy | Yes |